

Review Article

Secondary myopathy due to systemic diseases

Finsterer J, Löscher WN, Wanschitz J, Quasthoff S, Grisold W.
Secondary myopathy due to systemic diseases.

Acta Neurol Scand 2016; 134: 388–402.

© 2016 John Wiley & Sons A/S. Published by John Wiley & Sons Ltd.

Background – Some systemic diseases also affect the skeletal muscle to various degrees and with different manifestations. This review aimed at summarizing and discussing recent advances concerning the management of muscle disease in systemic diseases. **Method** – Literature review by search of MEDLINE, and Current Contents with appropriate search terms. **Results** – Secondary muscle disease occurs in infectious disease, endocrine disorders, metabolic disorders, immunological disease, vascular diseases, hematological disorders, and malignancies. Muscle manifestations in these categories include pathogen-caused myositis, muscle infarction, rhabdomyolysis, myasthenia, immune-mediated myositis, necrotising myopathy, or vasculitis-associated myopathy. Muscle affection may concern only a single muscle, a group of muscles, or the entire musculature. Severity of muscle affection may be transient or permanent, may be a minor part of or may dominate the clinical picture, or may be mild or severe, requiring invasive measures including artificial ventilation if the respiratory muscles are additionally involved. Diagnostic work-up is similar to that of primary myopathies by application of non-invasive and invasive techniques. Treatment of muscle involvement in systemic diseases is based on elimination of the underlying cause and supportive measures. The prognosis is usually fair if the causative disorder is effectively treatable but can be fatal in single cases if the entire musculature including the respiratory muscles is involved, in case of infection, or in case of severe rhabdomyolysis. **Conclusion** – Secondary muscle manifestations of systemic diseases must be addressed and appropriately managed. Prognosis of secondary muscle disease in systemic diseases is usually fair if the underlying condition is accessible to treatment.

**J. Finsterer¹, W. N. Löscher²,
J. Wanschitz², S. Quasthoff³,
W. Grisold⁴**

¹Krankenanstalt Rudolfstiftung, Vienna, Austria;

²Department of Neurology, Medical University of

Innsbruck, Innsbruck, Austria; ³Department of

Neurology, Graz Medical University, Graz, Austria;

⁴Department of Neurology, Kaiser-Franz-Josef Spital,
Vienna, Austria

Key words: system disease; endocrine disorders; infections; immunological disease; malignancy; muscle; myopathy; myasthenia; myopathy; rhabdomyolysis

J. Finsterer, Krankenanstalt Rudolfstiftung, Postfach 20,
1180 Vienna, Austria

Tel.: +43 1 71165 92085

Fax: +43 1 4781711

e-mail: fifigs1@yahoo.de

Accepted for publication February 9, 2016

Introduction

Systemic diseases are characterized by involvement or affection of the entire organism and contrast focal diseases, in which only a single organ or part of it is affected [1a]. Systemic diseases are also those affecting a system which is distributed all over the body, such as the nervous system (CNS), the vascular system, or the blood. Systemic diseases include for example leukemia, anemia, central nervous systemic diseases, vasculitis, diabetes,

thyroid dysfunction, hemochromatosis, rheumatological disorders, psoriasis, sarcoidosis, systemic lupus erythematosus, scleroderma, or mucoviscidosis (1). Some of these disorders secondarily affect the skeletal muscle, manifesting as myositis, muscle abscess, endocrine or metabolic myopathy, rhabdomyolysis, or myasthenia. The following review aims at summarizing and discussing recent advances concerning the management of muscle disease in systemic diseases, including infectious diseases, endocrinological disorders, metabolic

disorders, immunological disorders, vascular disorders, and hematological disorders.

Methods

Data for this review were identified by searches of MEDLINE, Current Contents, for references of relevant articles using the search terms 'infection', 'infectious', 'viral', 'bacterial', 'protozoal', 'helminthic', 'leukemia', 'anemia', 'diabetes', 'thyroid', 'hemochromatosis', 'rheumatological', 'psoriasis', 'lupus', 'scleroderma', 'sarcoidosis', 'mucoviscidosis', 'vasculitis', and 'vascular' in combination with 'skeletal muscle', 'myopathy', 'myositis', 'myasthenia', 'neuromuscular transmission', and 'rhabdomyolysis'. Randomized (blinded or open label) clinical trials, longitudinal studies, case series, and case reports were considered. Abstracts and reports from meetings were not included. Only articles published in English, French, Spanish, or German between 1966 and 2015 were included. Appropriate papers were studied and discussed for their suitability to be incorporated in this review. Due to limitations of space not all systemic diseases with muscle involvement could be discussed. Only the clinically most relevant were included.

Results

Classification of muscular manifestations of systemic diseases

Muscular manifestations in systemic diseases may be classified as acute, subacute, or chronic. Acute muscle manifestations include pathogen-caused myositis, muscle infarction, or rhabdomyolysis. Subacute or chronic muscular manifestations of systemic diseases include secondary endocrine or secondary metabolic myopathy, myasthenia, immune-mediated myositis, muscle abscess, or vasculitis with secondary myopathy (2). They may be further classified as transient or permanent, as mild or severe, as dominating or not dominating the clinical presentation, or as affecting a single muscle, a group of muscles, or the entire musculature.

Infectious diseases

Viral infections – Systemic viral infections may manifest in the muscle as myositis, rhabdomyolysis, or myasthenia. The most frequent among the muscle manifestations is myositis with a self-limiting course. More rarely, viral infections may manifest with rhabdomyolysis. Myasthenia due to viral infections is only reported in single cases.

Only few data about the clinical manifestations, frequency of muscle manifestations, and causative agents are available.

Myositis – In a retrospective study of 35 patients with viral myositis, muscle manifestations included localized pain of the calves (80%), lower limb weakness (71%), impaired ambulation (57%), and gait disturbance (40%) lasting on the average 3.6 days (3). Symptoms were associated with elevated muscle enzymes (3). Myositis may affect all muscles, a group of muscles (4, 5), or a single muscle (6). In a study of 355 pediatric patients with laboratory-confirmed influenza-B, 17.9% developed myositis (7). During the influenza pandemic in 2009, children developed more frequently myositis as compared to adults (8). Infections with human T-cell lymphotropic virus (HTLV-1) may manifest as axial myositis (4). Some of the viral infections may also cause orbital myositis, such as infections with the chickenpox virus (5). Rarely, a single muscle like the serratus anterior may be affected during flu (6).

Concerning the causative agents, viral myositis has been reported following infections with influenza-B (7), influenza-A (8), parainfluenza-1 (9), parvoviruses (10), HTLV-1 (11), Epstein-Barr virus (12), arboviruses (e.g., dengue myositis) (13), adenovirus (14), coxsackie (15), herpes (16), human immunodeficiency virus-1 (HIV-1) (17), or chickenpox (5). Most frequently myositis is due to infection with the influenza (7) or parainfluenza virus (9). These patients may present with acute onset calf pain, tenderness, or gait disturbance (9). Myositis in dengue infection can be fulminant and affect the respiratory muscles requiring artificial ventilation (18). Only rarely is myositis due to infection with parvovirus-B19 (10), HCV (19), or the West Nile virus (20). Chronic HCV infection may be also associated with dermatomyositis (21) or inclusion body myositis (22). Infection with the New Jersey polyomavirus NJPyV-2013 may cause vasculitic myopathy (2). Muscle-tropic viruses often spread to the CNS, which may dramatically increase morbidity and mortality (23). Whether patients with an underlying subclinical primary or secondary myopathy or those taking muscle-toxic drugs are particularly prone to develop viral myositis is unknown. Infection with HIV-1 may be associated with polymyositis, dermatomyositis, or inclusion body myositis (17). Polymyositis from HIV-1 may be complicated by the development of muscle abscesses (24). HIV-myositis may be even the initial manifestation of acquired immunodeficiency syndrome (AIDS) (25).

Rhabdomyolysis – Viral infections, particularly dengue and influenza-A, have been recognized as cause of rhabdomyolysis with considerable morbidity and mortality (26). Rhabdomyolysis as a complication of a viral infection occurs particularly in children and usually has a benign course (27). The main complication of rhabdomyolysis is acute renal failure, often requiring hemofiltration or hemodialysis (28). Rhabdomyolysis occurs in 1% of the patients with dengue infections. Risk factors for developing rhabdomyolysis include myalgia, arterial hypertension, and acute renal failure (29). More rarely rhabdomyolysis may be triggered by infections with influenza-A (30), influenza-B (31,32), parainfluenza (33), herpes-6 (34), varicella zoster (35), cytomegaly virus (CMV) (36), coronavirus-NL63 (37), chikungunya (38), Alkhurma hemorrhagic fever virus (39), or HIV-1 (40). Rhabdomyolysis in dengue fever infections can be life-threatening in some cases (41–43).

Myasthenia – Whether viral infections cause myasthenia is under debate. Well-known, however, is that infections may deteriorate myasthenia. There are, however, some reports indicating that a causal relation between a viral infection and the development of myasthenia may exist. In six Chinese patients with a West Nile virus infection, myasthenia developed 3–7 months after the infection (44). Further evidence for an association between viral infections and myasthenia are results of single-fiber EMG investigations in patients with influenza or echovirus infection showing a neuromuscular transmission defect (45).

Bacterial infections – Systemic bacterial infections manifest in the muscle as myositis or rhabdomyolysis.

Myositis – Bacterial infections associated with myositis usually manifest as bacterial polymyositis (46). Commonly, bacterial polymyositis is a purulent infection accompanied by muscle abscesses (pyomyositis) (46). The main etiologic agent of bacterial polymyositis is *Staphylococcus aureus* (46). Staphylococcal pyomyositis is a severe infection with high mortality being increasingly recognized in temperate climates (47). Pyomyositis may originate from a focal infection such as arthritis, sacroiliitis, a spinal abscess, or from bacteremia or sepsis. Pyomyositis may develop after trauma, may remain focal, and may resolve upon a nonsurgical approach (48). Pyomyositis may affect a single muscle, such as

the rectus femoris muscle (49), or may affect all muscles resulting in quadriplegia, as described in a patient after induction of chemotherapy for lymphoblastic leukemia (50). Pyomyositis may be diagnosed with ultrasound and culture of the aspirate (51). Pyomyositis of the iliopsoas muscle may be complicated by septic pulmonary embolism (47). These patients may require abscess drainage under CT-guidance (47). Presence of intramuscular hemangiomas seems to predispose for pyomyositis, as reported in a 4-year-old child with fatal meningitis (46). Sepsis from streptococcus group-G originating from arthritis may cause diffuse polymyositis without skin lesions or toxic shock syndrome (52). A severe form of bacterial myositis is streptococcal necrotizing myositis, which is often fatal (53). Muscle abscesses may also result from infection with *Klebsiella pneumoniae* (49) or *Mycobacterium tuberculosis* (54). A rare cause of systemic myositis may be infection with *Campylobacter jejuni* (55). Predisposing factors include skin penetration or impaired host immunocompetence (HIV-1, transplant recipient) (47).

Rhabdomyolysis – Occasionally, rhabdomyolysis may be a manifestation of a bacterial infection. Rhabdomyolysis has been particularly reported during infections with *Staphylococcus aureus* (56), *Salmonella* (57), *Brucella* (58), *Mycoplasma pneumoniae* (59), tuberculosis (60), tetanus (61), *Legionella* (62), or *Bacillus cereus* (63). In a 6-year-old girl, life-threatening rhabdomyolysis was triggered by streptococcus bovis sepsis (27).

Protozoal infections – Muscle manifestations of protozoal infections include myositis or rhabdomyolysis.

Myositis – Protozoal infections are a frequent cause of myositis. In some of these cases, myositis may be the dominant feature of the infection, such as in muscular sarcocystis (64). Causative agents include sarcocystis, plasmodium falciparum, toxoplasma gondii, neospora, microspora, borrelia, pleistophora, babesia, ehrlichia, or trypanosoma. Muscular sarcocystis is clinically characterized by myalgia with or without fever, and delayed onset of hyper-CKemia and eosinophilia with the possibility of relapses (65). Muscular sarcocystis is particularly prevalent in Malaysia (65). Other frequent protozoal infections with muscular involvement include toxoplasmosis (66) and malaria (67). In toxoplasmosis, severe polymyositis may be even the presenting manifestation (66). In a patient with AIDS, myositis due

to toxoplasmosis developed despite adequate antimicrobial treatment (68). Rarely, toxoplasmosis may be associated with dermatomyositis (69). Occasionally, falciparum malaria may manifest with myositis (67). A rare protozoic infection with muscle involvement is neosporosis (70). In immune-compromised patients, microsporidia, obligate intracellular parasites, may manifest as focal or generalized myositis (71,72). Focal myositis may be also a rare manifestation of Lyme disease (Lyme myositis) (73,74). Lyme myositis may even mimic dermatomyositis (75). Infection with *Borrelia burgdorferi* may also cause idiopathic inflammatory myopathy (76). In immune-compromised patients, myositis may be rarely caused by pleistophora (77). Ocular myositis may be caused by ehrlichiosis, babesiosis, or Lyme disease (78). In immune-compromised patients, ocular myositis may be also due to *Trypanosoma cruzi* infection (79).

Rhabdomyolysis – Rhabdomyolysis due to protozoal infections has been particularly reported in malaria (80). Another rare cause of rhabdomyolysis due to protozoal infection is babesiosis or ehrlichiosis (81). Rhabdomyolysis due to *Borrelia burgdorferi* infection has been reported only once (82).

Helminthic infections – Helminthic infestations are frequently associated with muscle disease. Helminthic infestations manifest in the muscle predominantly as myositis. Helminthes potentially affecting the muscle include *Toxocara* (toxocarosis) (83), *Echinococcus granulosus* (hydatidosis) (84), *Cysticercus* (cysticercosis) (85), *Trichinella* (trichinosis) (86), *Strongyloides* (strongyloidiasis) (87), *Haycocknema perplexum* (88), *Spirometra* (sparganosis) (89), *Fasciola* (fasciolosis) (90), or *Filaria* (filariasis). *Toxocara* infection may go along with lumbar myositis (83). In the tropics, visceral larva migrans (toxocarosis) may manifest as tropical pyomyositis requiring repeated debridement (91). Hydatid cysts from infestation with *Echinococcus* may rarely occur in a single muscle as the initial manifestation (84,92). Most commonly, liver and lung are affected (92). Hydatid cysts of the muscle have been occasionally observed in patients with primary muscle disease (93). Cysticercosis may initially manifest as ptosis if the lid elevator is affected (94). Cysticercosis may also manifest as ocular myositis (95). Focal cysticercal myositis may be diagnosed with muscle ultrasound or MRI (85). Trichinosis manifests clinically in the muscle as myalgias due to dermato-polymyositis (86, 96). *Trichinella* has a unique relation to the muscle as it is located

intracellularly. Patients may present with myalgia and fever, and elevated muscle enzymes (97). Rarely, trichinosis may go along with muscle weakness (98). In case of focal necrosis due to trichinosis, EMG may show profuse fibrillations (99). Later in the course, fibrosis and contractures may develop (99). *Strongyloides* rarely affects the musculature. Occasionally, patients taking steroids or immuno-suppressants may develop polymyositis from *strongyloides* infestation (87). In Australia, myositis may be due to infestation with the nematode *Haycocknema perplexum* (88). In single cases, sparganosis may manifest as ocular myositis (89). Rarely, cutaneous fascioliasis may cause myositis of the intercostal muscles (90). Filariasis rarely manifests as myositis with muscle swelling (100).

Endocrinological disorders

Diabetes – Diabetes is a catabolic condition which manifests in the muscle as diabetic myopathy (101). Diabetic myopathy encompasses a spectrum of abnormalities, including wasting, muscle inflammation, ischemia, infarction, hemorrhage, necrosis, fibrosis, or fatty atrophy (101). Clinical manifestations vary depending on the underlying abnormality. The most frequent muscle manifestation in diabetes is painless muscle wasting (diabetic amyotrophy), which is either due to a diabetic plexus lesion (102) or due to affection of satellite cells by the diabetes (103). A further frequent manifestation of diabetic myopathy is diabetic myonecrosis presenting as self-limiting condition with acute onset of swelling and severe muscle pain (104). Typically, patients have no fever, normal white blood cell count, normal blood sedimentation rate, but elevated C-reactive protein (104). Usually, myonecrosis occurs in poorly controlled diabetes (105). The diagnosis is established by muscle MRI, and the treatment of choice is bed rest and analgesics (104). An increasingly recognized manifestation of diabetic myopathy is diabetic muscle infarction, which is regarded as rare and occurs in long-standing diabetic patients (106). Clinical manifestations of muscle infarction include acute onset local pain, together with a focal, palpable mass lesion. The diagnosis is established by muscle MRI (106). Additionally, the expression level of Pax7, MyoD, myogenin, and fatal myosin-heavy-chain (MHC) is significantly decreased in diabetic myopathy (102).

Thyroid dysfunction – Hypothyroidism – Hypothyroidism is a well-known cause of muscle disease (hypothyroid myopathy) (107). In contrast to hyperthyroid myopathy, CK is usually elevated,

pain is frequent, and muscles can be swollen. Additionally, hypothyroidism due to autoimmune disease may be associated with dermatomyositis (108) or polymyositis (109, 110). Rarely, hypothyroidism may also go along with rhabdomyolysis (111). In infancy or childhood, hypothyroidism may manifest as Kocher–Debré–Semelaigne syndrome, characterized by lower limb or generalized muscle hypertrophy, myxedema, short stature, and cretinism. In adults, hypothyroidism may manifest as Hoffmann’s syndrome, characterized by muscle stiffness and muscle pseudo-hypertrophy. Muscle enzymes are generally elevated in hypothyroid myopathy. The EMG may show a myopathic, neuropathic, or a mixed pattern. Clinical manifestations of hypothyroid myopathy return to normal with hormone replacement therapy.

Hyperthyroidism – Graves’ disease may manifest in the skeletal muscle as mild and usually painless proximal weakness or as idiopathic ocular myositis (112). Myositis may respond favorably to thiazole without adding steroids (113). Thyrotoxicosis may go along with episodic muscle weakness due to polymyositis (113). Thyrotoxicosis may also manifest as acute or chronic bulbar muscle dysfunction (bulbar myopathy). If thyrotoxicosis leads to hypokalemia, generalized muscle weakness may ensue (thyrotoxic periodic paralysis). A rare manifestation of hyperthyroidism or thyrotoxicosis may be myokymia.

Hyper-/hypoparathyroidism – Hyperparathyroidism may cause muscle weakness (dropped head syndrome), muscle pain, or ischemic, calcifying myopathy (114, 115). Hyperparathyroidism may also go along with spontaneous rupture of the Achilles tendons. Rarely, the initial manifestation of hyperparathyroidism may be dysphagia. FDG-PET in hyperthyroidism may show tumors mimicking muscular metastases. Affection of the muscle in hypoparathyroidism may present as myopathy, neuromyotonia, or rhabdomyolysis (116).

Other endocrinopathies – Hypoadrenalism as well as hyperadrenalism may be complicated by generalized muscle weakness. In case of acromegaly, the muscles may be hypertrophic and stronger than normal, but later in the course proximal weakness may become evident.

Metabolic disorders

Various metabolic disorders secondarily affect the muscle. The most well-known are hemochromatosis, amyloidosis, and porphyria.

Hemochromatosis – Hereditary hemochromatosis and other iron-metabolism disorders resulting in iron overload may involve the skeletal muscle (iron-overload myopathy) which usually manifests as myalgias and fatigue (117, 118). Figures about the prevalence of iron-overload myopathy are highly variable. In a study of 46 patients with hereditary hemochromatosis, myopathy was diagnosed in none of them (119). In a study of 395 patients with hereditary hemochromatosis on the contrary, 43% were diagnosed with fibromyalgia (120). In a study of 88 patients with chronic fatigue syndrome, 2.6% had hereditary hemochromatosis (121). Among 10 patients under hemodialysis, muscle biopsy disclosed iron deposition in muscle fibers or macrophages in 70% of them (118). Clinically, these patients presented with proximal muscle weakness (118).

Amyloidosis – Muscle involvement is frequent in amyloidosis and manifests as amyloid myopathy (122), clinically characterized by muscle hypertrophy (muscle overgrowth), or weakness. Depending on the cause of amyloidosis, different types of amyloid (AA, AL) may be produced. Amyloidosis usually manifests systemically in all muscles but occasionally only a single muscle or a few muscles is/are affected (123). Amyloidosis is characterized by extracellular and perivascular deposition of AA or AL amyloid (122). Rarely, amyloid myopathy occurs in patients with hereditary transthyretin amyloidosis (ATTR) (124). Occasionally, systemic amyloidosis due to multiple myeloma may be exclusively detectable in the skeletal muscle as ring-fiber-like muscle fibers, staining positive for Congo-red (125). Rarely, amyloidosis due to IgD multiple myeloma may manifest as myositis (126). In systemic AL, amyloidosis amyloid myopathy may manifest with joint contractures (127). Amyloidosis of the muscle may also occur in monoclonal gammopathy (128). Histologically, amyloid myopathy can mimic inclusion body myopathy (129). In such cases, nonconophilic deposition of kappa-light chains can be seen as subsarcolemmal rings (128). Focal accumulations of amyloid may present as amyloidoma (tumoral amyloidosis) (130). Interstitial amyloid deposition in the muscle may rarely occur in patients with myopathy due to mutations in the anoctamin-5 gene (131).

Porphyrias – Porphyrias are diseases in which porphyrins, necessary to produce heme, accumulate. Most frequently, they manifest in the skin, brain, or peripheral nerves. More rarely, they manifest in the skeletal muscle. Acute intermittent

porphyria may go along with rhabdomyolysis (132). There are also reports about transient muscle weakness and muscle contractures in acute intermittent porphyria (133). Acute intermittent porphyria may also manifest as acute myalgias or muscle weakness (134).

Immunological disorders

A number of immunological disorders may involve the skeletal muscle presenting as polymyositis, dermatomyositis, rarely as inclusion body myositis, or as ocular myositis. The most important among these disorders are systemic lupus erythematosus (SLE), Sjögren syndrome, rheumatoid arthritis, systemic sclerosis, psoriasis, and the antisynthetase syndrome (ASS).

Systemic lupus erythematosus (SLE) – SLE is a chronic inflammatory, multisystem disease with a broad spectrum of clinical and serological abnormalities (135). SLE has been repeatedly reported to manifest with muscle disease. Muscle manifestations of SLE include polymyositis or rhabdomyolysis (Table 1). Some patients with SLE may even develop myositis together with rhabdomyolysis (136). Myositis is much more frequent than rhabdomyolysis. In a study of 15 SLE patients, 9% had developed clinical myositis (137). However, histopathological evidence of myositis was seen in 47% of these patients (137). Type-2 atrophy was the predominant histopathological finding. Occasionally, myositis may be the initial manifestation of SLE (138). Rarely, SLE may manifest as ocular myositis (139). Muscle involvement in SLE may also manifest as necrotizing autoimmune myopathy with muscle weakness and rhabdomyolysis (140). Polymyositis is also a feature of mixed connective tissue disease, including clinical and laboratory manifestations of SLE, scleroderma, and polymyositis along with high titers of anti-U1 and anti-U2-nRNP antibodies (141). Myositis may also occur in patients with SLE/polymyositis or SLE/dermatomyositis overlap syndrome (142). Rhabdomyolysis in SLE may be fatal in some cases (143).

Sjögren syndrome – The skeletal muscle is frequently affected in Sjögren syndrome usually manifesting as myalgia or weakness (144). In 3% of the patients, myositis has been described (144). In a study of 573 patients with Sjögren syndrome, myositis was found in 4.9% of them (145). Occasionally, muscle affection manifests as ocular myositis (146). Rarely, dermatomyositis may occur (147). Occasionally, Sjögren syndrome may

Table 1 Muscle manifestations of systemic diseases

Disorder	MP	FMS	PM	DM	IBM	MG	RM
Infectious disease							
Viral infections	+	+ (om)	+	+	+	+	+
Bacterial infections	–	+ (om)	+*	–	–	–	+
Protozoal infections	–	+ (om)	+	+	–	–	+
Helminthic infections	–	+ (om)	+	+	–	–	–
Endocrinological disorders							
Diabetes	+	–	–	–	–	–	–
Hypothyroid dysfunction	+	–	+	+	–	–	+
Hyperthyroid dysfunction	+	+ (om)	+	+	–	–	+
Hyperparathyroidism	+	–	–	–	–	–	–
Hypoparathyroidism	+	–	–	–	–	–	–
Hypo-/hyperadrenalism	+	–	–	–	–	–	–
Metabolic diseases							
Hemochromatosis	+	–	+	–	–	–	–
Amyloidosis	+	+	+	–	–	–	–
Porphyria	+	–	–	–	–	–	+
Immunological disorder							
Systemic lupus erythematosus	–	+ (om)	+	–	–	–	+
Sjögren syndrome	+	+ (om)	+	+	+	–	–
Rheumatoid arthritis	–	+	+	–	–	–	–
Systemic sclerosis	–	+	+	+	–	–	–
Psoriasis	+	+ (om)	+	+	–	–	–
Antisynthetase syndrome	–	+	+	–	–	–	–
Sarcoidosis			+		+		
Vascular diseases							
Behcet disease	–	+ (om)	+	–	–	–	–
Wegener	–r	+ (om)	–	–	–	–	–
Churg–Strauss syndrome	–	+ (om)	+	–	–	–	–
Hematological disease							
Sickle cell anemia	+	+	–	–	–	–	+
Neoplasms							
Leukemia	–	–	+	+	+	–	–
Lymphoma	–	+	+	+	–	–	–
Breast, lung, gastrointestinal	+*	–	–	–	–	–	–
Bladder tumor	+*	–	–	–	–	–	–

MP, myopathy; FMS, focal myositis; om, orbital myositis; PM, polymyositis. *Pyomyositis, abscess; IBM, inclusion body myositis; RM, rhabdomyolysis; necrotizing myopathy.

be also associated with polymyositis (148). In some patients, Sjögren syndrome was even associated with inclusion body myositis (149). The latter patients usually carried the HLA-DRB1 allele or its equivalent HLA-DR3 or the MHC ancestral haplotype. This is why the association of Sjögren syndrome and sporadic inclusion body myositis was assumed due to a genetic predisposition linked to MHC (150).

Rheumatoid arthritis – Affection of the skeletal muscles in rheumatoid arthritis manifests as rheumatoid myositis (151). Causes of rheumatoid arthritis are variable and include inflammation, drugs, impaired joint flexibility, or sedentarism (151). Muscle enzymes are usually highly elevated. Electromyography (EMG) may show short duration, low-amplitude, polyphasic motor unit action potentials (151). Active inflammation can be found on both muscle ultrasound and MRI

(151). Muscle biopsy may show nonspecific findings, such as changes in fiber size or internal structure, pleomorphic mitochondria, dilated sarcolemmal nuclei, a trend toward type-II-fibers, fiber atrophy, degenerative/regenerative modifications, and inflammatory features such as patchy B- or T-cell infiltrates, mainly perivascularly or endomysially, but also in the perimysial compartment (151). In rare cases, rheumatoid myositis may be complicated by compartment syndrome (152) or may affect the extraocular muscles (ocular myositis) (153, 154). Adalimumab has been reported to induce myositis in rheumatoid arthritis. Eosinophilic myositis has been reported in a patient with myotonic dystrophy-1 and rheumatoid arthritis (155).

Systemic sclerosis – Systemic sclerosis is characterized by arthralgia, synovitis, contractures, tendon friction rubs, tenosynovitis, and muscle disease (156). Muscle involvement in systemic sclerosis is frequent (156). Muscular manifestations of systemic sclerosis include myositis with myalgia and weakness (156). Rarely, patients with systemic sclerosis may develop inclusion body myositis (157). In a study of 1145 patients with systemic sclerosis, 5.6% had elevated CK. This subset of patients had a poor prognosis impacting survival, particularly in male with early onset, topol and RNP autoantibodies, and interstitial lung disease (158). If muscle weakness includes the head extensor muscles, dropped head syndrome may ensue (159). Muscle MRI is an appropriate tool to support the diagnosis of muscle involvement in systemic sclerosis (160). In some patients, myositis responded favorably to immunoglobulin (161).

Psoriasis – Psoriasis is a noninfectious, inflammatory dermatological disorder, which presents as systemic disease if joints, bands, eyes, arteries, or the heart are additionally involved. Psoriasis is associated with an increased risk of diabetes and ischemic stroke. Only rarely muscle involvement has been reported. In a study of 11370 patients with psoriasis, 13 had a pathologically confirmed myopathy (0.11%) (162). Eleven had generalized inflammatory myopathy and 2 had focal muscular inflammation (162). In two-thirds of these patients, onset of psoriasis preceded that of myopathy. Patients receiving a TNF-alpha blocker had an increased risk of developing myositis (162). Rarely, orbital myositis has been reported (163). FDG-PET may show an increased tracer-uptake in affected muscles (164). In a single case, psoriasis was associated with dermatomyositis and antibodies against the glomerular basement membrane (165).

Antisynthetase syndrome (ASS) – ASS is due to auto-antibodies directed against aminoacyl-transfer ribonucleic acid (tRNA) synthase enzymes such as anti-Jo1, anti-PL7, or anti-PL12 (166). Clinical manifestations include fever, nonerosive polyarthritides, Raynaud phenomenon, mechanic's hands, myocarditis (42% of patients), interstitial lung disease, which is frequently associated with anti-Jo1 antibodies, pulmonary hypertension, progressive multifocal leucoencephalopathy, or myositis with myalgias (166, 167). Presence of interstitial lung disease is a major prognostic factor (168). Treatment of choice is the application of corticosteroids with or without immunosuppressive agents (168).

Vascular disorders

Muscle function not only depends on appropriate innervation and energy production, but also on sufficient blood perfusion. Muscle perfusion may depend not only on cardiac function but also on muscle artery contractility. Physiologically, endothelial cells produce basal and stimulated nitric oxide (NO). During exercise, NO production is stimulated, which contributes to exercise-induced muscle hyperemia. In patients with reduced NO production due to reduced activity of NO-synthetase (NOS), reduced microcirculation contributes to exercise-induced muscle fatigue. NO deficiency results in muscle hypoperfusion with decreased provision of nutrients and thus decreased protein production (169). Microvascular perfusion is particularly compromised in systemic vasculitis, which includes Behcet disease, Wegener's granulomatosis, and Churg–Strauss syndrome.

Behcet disease – Behcet disease is a form of systemic vasculitis with the classical triad of oral ulcers, genital ulcers, and uveitis (152). Involvement of the skeletal muscle is rare (152). The most common muscle manifestation is myositis. Myositis usually shows a focal distribution (170). In single cases, orbital myositis has been reported (171). Only a single extra-ocular muscle may be affected (172). Only rarely may myositis dominate the clinical presentation (173). Development of myositis in Behcet disease may be triggered by stress such as surgery (174). Occasionally, myositis may be of the necrotizing type (175). Generalized myositis may be diagnosed by muscle ultrasound, CT, or MRI (152). Generalized

myositis in Behcet disease may occasionally respond favorably to cyclosporine (176).

Wegener's granulomatosis – Wegener's granulomatosis (granulomatosis with polyangiitis) is a systemic necrotizing vasculitis which is associated with granulomatous infection of the nasopharynx, the sinuses, the oropharynx, and the lower bronchial airways. Muscle involvement is infrequent and usually manifests as myositis. The most common type of myositis in Wegener's granulomatosis is ocular myositis (177). If myositis predominantly affects the lower limb muscles, it may go along with muscle weakness and gait disturbance (178).

Churg–Strauss syndrome – Churg–Strauss syndrome, also known as eosinophilic granulomatosis with polyangiitis, is a granulomatous vasculitis of the small arteries accompanied by infiltrates of eosinophils. It manifests clinically in three stages, initially as allergic cold and asthma, followed by eosinophilic infiltration of the lung and intestines and systemic vasculitis. Myositis is a rare muscle manifestation of the disease and presents with myalgia, fever, and muscle weakness (179). Myositis in Churg–Strauss syndrome may not only concern all muscles resulting in polymyositis (180) but may occur focally as orbital myositis (181).

Hematological disorders

Hematological disorders are rarely associated with muscle disease. Muscle involvement has been particularly reported in sickle cell anemia (182). Muscle affection in sickle cell anemia includes myalgia, focal myopathy, focal myositis, pyomyositis, myonecrosis, fibrosis, fasciitis, or rhabdomyolysis. Muscle involvement is more frequent in hematological neoplasms, but they are described in more detail below.

Neoplasms

Muscle disease in neoplasms is a paraneoplastic phenomenon and includes focal or generalized myositis, polymyositis, dermatomyositis, or necrotizing myopathy. Neoplasms associated with muscle disease include leukemia, lymphomas, or other solid tumors.

Leukemia – Polymyositis/dermatomyositis are symmetric, proximal, paraneoplastic, inflammatory myopathies with or without distinct cutaneous eruptions (183). They have been long

recognized in association with cancer (183). Only rarely may polymyositis/dermatomyositis be associated with acute myelocytic leukemia (183). In single cases, chronic lymphatic leukemia may go along with inclusion body myositis (184, 185). Pyomyositis may be the initial presentation not only of chronic myeloid leukemia (186) but also of acute lymphocytic leukemia (187). In a girl with secondary acute myelogenous leukemia following chemotherapy, tuberculous myositis developed (188). Chemotherapy for leukemia may occasionally induce pyomyositis (189).

Lymphoma – Lymphoma is frequently associated with muscle disease, particularly with polymyositis or dermatomyositis (190). B-cell lymphoma, T-cell-lymphoma, and Hodgkin's lymphoma have been reported in association with dermatomyositis or polymyositis (191). In a study of 32 patients with polymyositis/dermatomyositis, 20 had B-cell lymphoma, four had T-cell lymphoma, and two had Hodgkin's lymphoma (191). In single cases, B-cell lymphoma manifested with isolated myositis of a single extra-ocular muscle (192). Non-Hodgkin lymphoma may directly develop inside the muscle.

Other malignancies – Paraneoplastic myopathy has been reported also in a number of other neoplasms. Lung, gastrointestinal, and breast carcinomas are frequently associated with necrotizing myopathy. The bladder transitional cell tumor may cause necrotizing myopathy with pipestem capillaries. Waldenström's macroglobulinemia may go along with antidecorin (BJ) myopathy. Patients with thymoma may develop rippling muscle syndrome. Patients with paraproteinemia (M-protein, $\kappa > \lambda$ light chains, IgG) or carcinoids may present with scleromyxedema.

Diagnosis

Methods to diagnose muscle manifestations of systemic disease are the same as those applied for diagnosing primary muscle disease. The basis is a thorough individual and family history and a thorough clinical exam. Determination of muscle enzymes, EMG, muscle imaging, and a muscle biopsy may be of additional help. FDG-PET may show increased muscular tracer-uptake in myositis (164) or tumors. Viral infections causing myositis may be diagnosed by detection of serum antibodies against viruses or by PCR. CK values may be higher during the acute stage of an influenza infection than during

the convalescence stage (193). Determination of various muscle-specific antibodies or auto-antibodies, such as anti-Jo1, anti-PL7, anti-PL12 (ASS) (166), anti-EJ, anti-OJ, anti-SRP, anti-Mi-2, anti-PM-Scl75, anti-PM-Scl100, and anti-Ku (overlap syndromes) may be necessary to establish the diagnosis of muscle disease in immunological disorders. U1-nRNP antibodies may be determined when suspecting SLE, scleroderma, or polymyositis overlap syndrome. Topo1 and RNP antibodies may be positive in myopathy from systemic sclerosis (158). Antidecorin antibodies (BJ antigen) may indicate Waldenström's macroglobulinemia. Determination of ryanodine-receptor antibodies may be helpful for diagnosing myasthenia gravis or myositis, and determination of monoclonal antibodies (M-proteins) may suggest scleromyxedema. Single-fiber EMG may show a disturbed neuromuscular transmission during the acute stage of influenza or echovirus infections (45). Disturbed neuromuscular transmission may explain muscle weakness and fatigue during a viral infection (45). If muscle imaging reveals an enhancing lesion with a fluid density and needle aspiration shows pus, *Staphylococcus aureus* is growing in 85% of the cases (194). Before diagnosing a secondary myopathy, a primary myopathy needs to be excluded (195).

Treatment

Treatment of muscle involvement in systemic diseases is mainly based on the treatment of the underlying disorder. Additionally, symptomatic measures for pain, muscle cramps, muscle stiffness, can be applied. Symptomatic measures for myositis may also include nonsteroidal analgesic drugs, steroids, immunoglobulin, or immunosuppressants. Diabetic myonecrosis responds favorably to bed rest and analgesics. In case of immune-mediated myasthenia, cholinergic drugs, steroids, or immuno-suppressants may be necessary. In case of vasculitis-related myopathy, immuno-suppression may be beneficial. Infectious myositis may respond to adequate antibiotic treatment. Helminthic infections may respond to antihelminthics with or without steroids. In case of abscess formation, puncture and drainage or resection may be indicated. In severe pyomyositis due to toxocarosis, repeated debridement may be inevitable (91). In a rare case of myopathy associated with Whipple disease antibiotics exhibited a beneficial effect on muscle manifestations (196). In case of severe rhabdomyolysis with renal insufficiency diuretics, hemofiltration or hemodialysis

may beneficially influence the muscle pathology. Most cases of muscle involvement in systemic disease profit from physiotherapy.

Limitations

Systemic disease is not addressed in this review because of limited space, to few reports in the literature, or low frequency of muscle involvement, include aspergillosis, celiac disease (197), Henoch-Schoenlein purpura, Crohn's disease, mucoviscidosis, sarcoidosis, AMPA-associated immune encephalitis (198), renal myopathy, and vitamin D deficiency (199).

Clinical implications and summary

For treating physician, it is essential to know about muscular involvement in infectious, endocrine, metabolic, immunogenic, vascular, hematological diseases, or neoplasms. As soon as muscle involvement is suspected, referral to the neurologist is inevitable, and appropriate diagnostic measures as outlined above need to be initiated. In emergency cases due to renal or respiratory compromise, the treating neurologist must instantly manage and supervise the diagnostic procedures to initiate appropriate treatment in due time. In case of chronic muscle involvement, diagnostic steps may be taken more slowly but may be more invasive including abscess puncture or muscle biopsy. Particularly in infectious diseases, it is important to precisely determine the causative agent to apply the most specific antimicrobial agents with the highest effect. Muscle involvement in systemic diseases needs to be recognized and thoroughly investigated, as some cases may take a rapid or fulminant course with a high probability of an unfavorable or even fatal outcome.

Acknowledgment

None.

Conflict of interest

The authors have nothing to declare.

References

1. DORLAND. Dorland's Illustrated Medical Dictionary, 32nd edn. Philadelphia: Saunders, 2011.
2. MISHRA N, PEREIRA M, RHODES RH et al. Identification of a novel polyomavirus in a pancreatic transplant recipient with retinal blindness and vasculitic myopathy. *J Infect Dis* 2014;**210**:1595–9.

3. CARDIN SP, MARTIN JG, SAAD-MAGALHÃES C. Clinical and laboratory description of a series of cases of acute viral myositis. *J Pediatr (Rio J)* 2015;**91**:442–7.
4. MATSUURA E, YOSHIMURA A, NOZUMA S, HIGUCHI I, KUBOTA R, TAKASHIMA H. Clinical presentation of axial myopathy in two siblings with HTLV-1 associated myelopathy/tropical spastic paraparesis (HAM/TSP). *BMC Neurol* 2015;**15**:18.
5. KIM JH, LEE SJ, KIM M. External ophthalmoplegia with orbital myositis in an adult patient after chickenpox infection. *BMJ Case Rep* 2014. pii: bcr2013202415. doi: 10.1136/bcr-2013-202415
6. DANDOLO A, BANERJEE A. Benign acute myositis in a 17-year-old boy. *Arch Pediatr* 2013;**20**:779–82.
7. MOON JH, NA JY, KIM JH et al. Neurological and muscular manifestations associated with influenza B infection in children. *Pediatr Neurol* 2013;**49**:97–101.
8. REDLBERGER-FRITZ M, HIRK S, BUCHINGER D et al. Distinct differences in clinical manifestation and viral laboratory parameters between children and adults with influenza A(H1N1)pdm09 infection—a retrospective comparative analysis. *J Med Virol* 2014;**86**:1048–55.
9. TIPPETT E, CLARK R. Benign acute childhood myositis following human parainfluenza virus type-1 infection. *Emerg Med Australas* 2013;**25**:248–51.
10. ÇAKIRCA M, KARATOPRAK C, UGURLU S, ZORLU M, KISKAÇ M, ÇETIN G. Parvovirus B19 infection as a cause of acute myositis in an adult. *Rev Bras Reumatol* 2015;**55**:185–8.
11. GILBERT DT, MORGAN O, SMIKLE MF, SIMEON D, BARTON EN. HTLV-1 associated polymyositis in Jamaica. *Acta Neurol Scand* 2001;**104**:101–4.
12. LAMABADUSURIYA SP, WITHARANA N, PREETHIMALA LD. Viral myositis caused by Epstein-Barr virus (EB virus) in children. *Ceylon Med J* 2002;**47**:38.
13. VERMA R, SAHU R, HOLLA V. Neurological manifestations of dengue infection: a review. *J Neurol Sci* 2014;**346**:26–34.
14. SAKATA H, TAKETAZU G, NAGAYA K et al. Outbreak of severe infection due to adenovirus type 7 in a paediatric ward in Japan. *J Hosp Infect* 1998;**39**:207–11.
15. GÓMEZ RODRÍGUEZ N, IBÁÑEZ RUÁN J, GONZÁLEZ RODRÍGUEZ M. Coxsackie virus infection associated with myositis and polyarthrits. *An Med Interna* 2008;**25**:90–2.
16. KIM HT, MOON SY, LEE KH. Acute orbital myositis before Herpes zoster ophthalmicus. *Korean J Anesthesiol* 2012;**62**:295–6.
17. ROEDLING S, PEARL D, MANJI H, HANNA MG, HOLTON JL, MILLER RF. Unusual muscle disease in HIV infected patients. *Sex Transm Infect* 2004;**80**:315–7.
18. PALIWAL VK, GARG RK, JUYAL R et al. Acute dengue virus myositis: a report of seven patients of varying clinical severity including two cases with severe fulminant myositis. *J Neurol Sci* 2011;**300**:14–8.
19. AGHA B, RASHEED M, KENYON L, JIMENEZ SA. Hepatitis C virus infection, inflammatory myopathy, and pulmonary fibrosis: are they related? *J Clin Rheumatol* 2002;**8**:44–9.
20. SMITH RD, KONOPLEV S, DECOURTEN-MYERS G, BROWN T. West Nile virus encephalitis with myositis and orchitis. *Hum Pathol* 2004;**35**:254–8.
21. KEE KM, WANG JH, LEE CM, CHANGCHIEN CS, ENG HL. Chronic hepatitis C virus infection associated with dermatomyositis and hepatocellular carcinoma. *Chang Gung Med J* 2004;**27**:834–9.
22. TSURUTA Y, YAMADA T, YOSHIMURA T et al. Inclusion body myositis associated with hepatitis C virus infection. *Fukuoka Igaku Zasshi* 2001;**92**:370–6.
23. WATSON NB, SCHNEIDER KM, MASSA PT. SHP-1-dependent macrophage differentiation exacerbates virus-induced myositis. *J Immunol* 2015;**194**:2796–809.
24. WIDROW CA, KELLIE SM, SALTZMAN BR, MATHUR-WAGH U. Pyomyositis in patients with the human immunodeficiency virus: an unusual form of disseminated bacterial infection. *Am J Med* 1991;**91**:129–36.
25. DE LUIS A, MONTALBÁN J, FERNÁNDEZ JM et al. Polymyositis as the first manifestation of acquired immunodeficiency syndrome. *Rev Clin Esp* 1991;**188**:249–51.
26. FADILA MF, WOOL KJ. Rhabdomyolysis secondary to influenza a infection: a case report and review of the literature. *N Am J Med Sci* 2015;**7**:122–4.
27. PAN HH, CHANG HM, CHEN SM, HUNG TW, LUE KH, SHEU JN. Life-threatening rhabdomyolysis induced by *Streptococcus bovis* sepsis in a 6-year-old girl presented to the pediatric emergency department. *Pediatr Emerg Care* 2012;**28**:469–71.
28. CALLADO RB, CARNEIRO TG, PARAHYBA CC. Lima Nde A, da Silva Junior GB, Daher Ede F. Rhabdomyolysis secondary to influenza A H1N1 vaccine resulting in acute kidney injury. *Travel Med Infect Dis* 2013;**11**:130–3.
29. HUANG SY, LEE IK, LIU JW, KUNG CT, WANG L. Clinical features of and risk factors for rhabdomyolysis among adult patients with dengue virus infection. *Am J Trop Med Hyg* 2015;**92**:75–81.
30. SATO E, NAKAMURA T, KOIDE H. Rhabdomyolysis induced by influenza A infection: case report and review of literature. *Ther Apher Dial* 2011;**15**:208–9.
31. FRÖHLICH G, GRAWE A, SCHMIDT N, ROTH R, MÄRKER-HERMANN E. Difficult course of rhabdomyolysis in influenza A/H1N1. *Dtsch Med Wochenschr* 2013;**138**:2351–4.
32. GHOSH JB. Influenza-B associated rhabdomyolysis with acute renal failure. *Indian Pediatr* 2013;**50**:1160–1.
33. DOUVOYIANNIS M, KIELBASA JM, CHANDRASEKHARAN GM, HOLMES CL, GOMEZ MR. Rhabdomyolysis associated with parainfluenza virus. *Case Rep Infect Dis* 2013;**2013**:650965.
34. FUJINO M, OHASHI M, TANAKA K, KATO T, ASANO Y, YOSHIKAWA T. Rhabdomyolysis in an infant with primary human herpesvirus 6 infection. *Pediatr Infect Dis J* 2012;**31**:1202–3.
35. MARTÍNEZ-MARTÍNEZ ML, CÓRDOBA-SORIANO JG, CALBO-MAYO J, MELERO BASCONES M. Varicella complicated by rhabdomyolysis. *Actas Dermosifiliogr* 2013;**104**:448–9.
36. JUNG HY, KIM KH, PARK SC et al. Rhabdomyolysis associated with cytomegalovirus infection in kidney transplant recipients. *Transpl Infect Dis* 2014;**16**:993–8.
37. CABEÇA TK, BELLEI N. Human coronavirus NL-63 infection in a Brazilian patient suspected of H1N1 2009 influenza infection: description of a fatal case. *J Clin Virol* 2012;**53**:82–4.
38. SAM IC, KAMARULZAMAN A, ONG GS et al. Chikungunya virus-associated death in Malaysia. *Trop Biomed* 2010;**27**:343–7.
39. RAVANINI P, HASU E, HUHTAMO E et al. Rhabdomyolysis and severe muscular weakness in a traveler diagnosed with Alkhurma hemorrhagic fever virus infection. *J Clin Virol* 2011;**52**:254–6.

40. HALPERIN J, RICHEY LE, MUSHATT D. The modern day masquerader: a rare case of acute HIV infection with neurologic complications and a review of the literature. *J La State Med Soc* 2012;**164**:76–80.
41. GARG RK, MALHOTRA HS, JAIN A, MALHOTRA KP. Dengue-associated neuromuscular complications. *Neurol India* 2015;**63**:497–516.
42. VERDOLIN LD, BORNER AR, MUSSI H, GISMONDI RA, SCHAU B, RAMOS RC. Rhabdomyolysis associated with dengue fever in a lupic patient. *Rev Bras Reumatol* 2014;**54**:318–21.
43. UTHAMALINGAM P, NADA R, RAMACHANDRAN R, RAYAT CS, SAKHUJA V, VASHISTA RK. Acute renal failure due to Dengue myositis: a rare cause of pigment cast nephropathy. *Clin Kidney J* 2013;**6**:662–3.
44. LEIS AA, SZATMARY G, ROSS MA, STOKIC DS. West Nile virus infection and myasthenia gravis. *Muscle Nerve* 2014;**49**:26–9.
45. FRIMAN G, SCHILLER HH, SCHWARTZ M. Disturbed neuromuscular transmission in viral infections. *Scand J Infect Dis* 1977;**9**:99–103.
46. OLIVEIRA CC, MARQUES ME. Meningitis and pneumococcal pyomyositis in a child with intramuscular hemangiomas: an autopsy case report. *Int J Clin Exp Pathol* 2014;**7**:4523–7.
47. SAVVIDOU S, KALOGIANNIS E, TSAKIRI K, GAVRA M, TSONA A. Primary pyomyositis and disseminated septic pulmonary emboli: a reactivated staphylococcal infection? *Braz J Infect Dis* 2014;**18**:457–61.
48. KARAKATSANIS A, LYKODIS P, NASTOS C et al. Bilateral obturator pyomyositis with visceral involvement in an immunocompetent adult patient treated without surgery. *Surg Infect (Larchmt)* 2014;**15**:651–5.
49. LIAO WH, LAI CC, HUANG SH, LIN SH. Rectus femoris pyomyositis caused by *Klebsiella pneumoniae*. *Surg Infect (Larchmt)* 2014;**15**:464–5.
50. MONTAZERI N, ATHALE UH, FULFORD M, TARNOPOLSKY MA. Pyomyositis Causing Temporary Quadriplegia During Induction Therapy for Acute Lymphoblastic Leukemia: Case Report and Review of the Literature. *J Pediatr Hematol Oncol* 2015;**37**:223–6.
51. KUMAR MP, SEIF D, PERERA P, MAILHOT T. Point-of-care ultrasound in diagnosing pyomyositis: a report of three cases. *J Emerg Med* 2014;**47**:420–6.
52. DENG W, FARRICIELLI L. Group G streptococcal sepsis, septic arthritis and myositis in a patient with severe oral ulcerations. *BMJ Case Rep* 2014. pii: bcr2013200338. doi: 10.1136/bcr-2013-200338.
53. HOURMOZDI JJ, HAWLEY DA, HADI CM, TAHIR B, SEUPAUL RA. Streptococcal necrotizing myositis: a case report and clinical review. *J Emerg Med* 2014;**46**:436–42.
54. LAI KL, SHI ZY, CHAO WC, LIU PY, WU LH. Tuberculous muscle abscess. *QJM* 2014;**107**:683–4.
55. FUJIKI Y, KOTANI T, TAKEUCHI T, MAKINO S. Systemic myositis due to *Campylobacter* infection. *Scand J Rheumatol* 2014;**43**:78–80.
56. RAVRY C, FEDOU AL, DUBOS M et al. Severe rhabdomyolysis associated with staphylococcus aureus acute endocarditis requiring surgery. *Surg Infect (Larchmt)* 2015;**16**:840–2.
57. AL SHAMKHANI W, AJAZ Y, SAEED JAFAR N, ROY NARAYANAN S. Myocarditis and rhabdomyolysis in a healthy young man caused by salmonella gastroenteritis. *Case Rep Infect Dis* 2015;**2015**:954905.
58. NAHA K, KARANTH S, DASARI S, PRABHU M. Polymyositis-like syndrome with rhabdomyolysis in association with brucellosis. *Asian Pac J Trop Med* 2012;**5**:755–6.
59. CONSILVIO NP, RAPINO D, SCAPARROTTA A et al. Mycoplasma pneumoniae infection with rhabdomyolysis in a child. *Infez Med* 2014;**22**:48–50.
60. TEDESCO A, GUGLIEMETTI L, CONTI M, CAZZADORI A, CONCIA E. Pulmonary tuberculosis mimicking a severe community-acquired pneumonia with rhabdomyolysis. *Infez Med* 2013;**21**:146–8.
61. EL KORAICHI A, AGGOUG B, TADILI J, BENJELLOUN MY, EL HADDOURY M, EL KETTANI SE. Severe tetanus complicated with rhabdomyolysis and acute renal failure: a pediatric case report. *Arch Pediatr* 2012;**19**:726–8.
62. SHIMURA C, SARAYA T, WADA H et al. Pathological evidence of rhabdomyolysis-induced acute tubulointerstitial nephritis accompanying *Legionella pneumophila* pneumonia. *J Clin Pathol* 2008;**61**:1062–3.
63. GÓMEZ-HERRUZ P, GIL-FERNÁNDEZ JJ, GUILLEN H, ARIZCORRETA A. Fatal *Bacillus cereus* sepsis and rhabdomyolysis in a patient with Hodgkin's disease. *Enferm Infecc Microbiol Clin* 2011;**29**:232–3.
64. FAYER R, ESPOSITO DH, DUBEY JP. Human infections with *Sarcocystis* species. *Clin Microbiol Rev* 2015;**28**:295–311.
65. ESPOSITO DH, STICH A, EPELBOIN L et al. , Tioman Island Sarcocystosis Investigation Team. Acute muscular sarcocystosis: an international investigation among ill travelers returning from Tioman Island, Malaysia, 2011–2012. *Clin Infect Dis* 2014;**59**:1401–10.
66. CUOMO G, D'ABROSCA V, RIZZO V et al. Severe polymyositis due to *Toxoplasma gondii* in an adult immunocompetent patient: a case report and review of the literature. *Infection* 2013;**41**:859–62.
67. SINNAH R, LYE W. Acute renal failure from myoglobinuria secondary to myositis from severe falciparum malaria. *Am J Nephrol* 2000;**20**:339–43.
68. KUNG DH, HUBENTHAL EA, KWAN JY, SHELBURNE SA, GOODMAN JC, KASS JS. Toxoplasmosis myelopathy and myopathy in an AIDS patient: a case of immune reconstitution inflammatory syndrome? *Neurologist* 2011;**17**:49–51.
69. DA SILVA BS, PEREIRA RM, DE CARVALHO JF. Juvenile dermatomyositis and toxoplasmosis: a rare association. *Acta Reumatol Port* 2009;**34**:546–50.
70. LONGSHORE RC. What is your neurologic diagnosis? *Neospora caninum*-induced myositis. *J Am Vet Med Assoc* 1996;**208**:667–8.
71. FIELD AS, PAIK JY, STARK D et al. Myositis due to the microsporidian *Anncalia* (*Brachiola*) *algerae* in a lung transplant recipient. *Transpl Infect Dis* 2012;**14**:169–76.
72. PATEL AK, PATEL KK, CHICKABASAVIAH YT et al. Microsporidial polymyositis in human immunodeficiency virus-infected patients, a rare life-threatening opportunistic infection: clinical suspicion, diagnosis, and management in resource-limited settings. *Muscle Nerve* 2015;**51**:775–80.
73. HOLMGREN AR, MATTESON EL. Lyme myositis. *Arthritis Rheum* 2006;**54**:2697–700.
74. SAUER A, SPEEG-SCHATZ C, HANSMANN Y. Two cases of orbital myositis as a rare feature of Lyme borreliosis. *Case Rep Infect Dis* 2011;**2011**:372470.
75. WATON J, PINAULT AL, POUAHA J, TRUCHETET F. Lyme disease could mimic dermatomyositis. *Rev Med Interne* 2007;**28**:343–5.
76. GAN L, MILLER FW. State of the art: what we know about infectious agents and myositis. *Curr Opin Rheumatol* 2011;**23**:585–94.
77. GRAU A, VALLS ME, WILLIAMS JE, ELLIS DS, MUNTANÉ MJ, NADAL C. Myositis caused by *Pleistophora* in a patient with AIDS. *Med Clin (Barc)* 1996;**107**:779–81.

78. PENDSE S, BILYK JR, LEE MS. The ticking time bomb. *Surv Ophthalmol* 2006;**51**:274–9.
79. DOS SANTOS SDE S, ALMEIDA GM, MONTEIRO ML et al. Ocular myositis and diffuse meningoencephalitis from *Trypanosoma cruzi* in an AIDS patient. *Trans R Soc Trop Med Hyg* 1999;**93**:535–6.
80. PRABHAKAR KS, RATHORE SS, CHOUDHURY TA, KISHAN A, GUPTA T, PRAKASH J. Rhabdomyolysis induced acute renal failure: a rare complication of falciparum malaria. *J Assoc Physicians India* 2014;**62**:865–6.
81. JAVED MZ, SRIVASTAVA M, ZHANG S, KANDATHIL M. Concurrent babesiosis and ehrlichiosis in an elderly host. *Mayo Clin Proc* 2001;**76**:563–5.
82. JEANDEL C, PERRET C, BLAIN H, JOUANNY P, PENIN F, LAURAIN MC. Rhabdomyolysis with acute renal failure due to *Borrelia burgdorferi*. *J Intern Med* 1994;**235**:191–2.
83. BELLANGER AP, RUNGE M, WENDLING D, HUMBERT P. Lumbar myositis associated with *Toxocara* spp. infection. *Reumatol Clin* 2014;**10**:54–5.
84. BOYACI N, BOYACI A, KARAKAS E, ALTAY MA. Primer hydatid cyst of the deltoid muscle. *BMJ Case Rep* 2013. pii: bcr2013200209. doi: 10.1136/bcr-2013-200209.
85. TRIPATHY SK, SEN RK, AKKINA N, HAMPANNAVAR A, TAHASILDAR N, LIMAYE R. Role of ultrasonography and magnetic resonance imaging in the diagnosis of intramuscular cysticercosis. *Skeletal Radiol* 2012;**41**:1061–6.
86. DUBEY ML, KHURANA S, SINGHAL L, DOGRA S, SINGH S. Atypical trichinellosis without eosinophilia associated with osteomyelitis. *Trop Doct* 2011;**41**:244–6.
87. McNEELY DJ, INOUE T, TAM PY, RIPLEY SD. Acute respiratory failure due to strongyloidiasis in polymyositis. *J Rheumatol* 1980;**7**:745–50.
88. BASUROY R, PENNISI R, ROBERTSON T et al. Parasitic myositis in tropical Australia. *Med J Aust* 2008;**188**:254–6.
89. KUBOTA T, ITOH M. Sparganosis associated with orbital myositis. *Jpn J Ophthalmol* 2007;**51**:311–2.
90. LE XUAN T, HUNG NT, WAIKAGUL J. Cutaneous fascioliasis: a case report in Vietnam. *Am J Trop Med Hyg* 2005;**72**:508–9.
91. LAMBERTUCCI JR, RAYES A, SERUFO JC, TEIXEIRA DM, GERSPACHER-LARA R, NASCIMENTO E. Brasileiro Filho G, Silva AC. Visceral larva migrans and tropical pyomyositis: a case report. *Rev Inst Med Trop Sao Paulo* 1998;**40**:383–5.
92. NOTARNICOLA A, MORETTI L, PANELLA A, MARGARI AG, CIMINO A, PESCE V, MORETTI B. Case report of a primary multiloculate muscular cystic hydatidosis. *Chir Organi Mov* 2009;**93**:79–83.
93. SCHIMRIGK K, EMSER W. Parasitic myositis by *Echinococcus alveolaris*. Report of a family with myotonia congenita. *Eur Neurol* 1978;**17**:1–7.
94. KUNDR A, KUNDR SN. Unilateral ptosis due to cysticercosis of extraocular muscle. *Indian J Pediatr* 2004;**71**:181–2.
95. SEKHAR GC, LEMKE BN. Orbital cysticercosis. *Ophthalmology* 1997;**104**:1599–604.
96. CERVERA-CASTILLO H, TORRES-CABALLERO V, MARTÍNEZ-GARCÍA E, BLANCO-FAVELA F. Human trichinosis. A case simulating polymyositis. *Rev Med Inst Mex Seguro Soc* 2009;**47**:323–6.
97. LEVY Y, RIMBROT S, RAZ R. Myalgia, fever, abnormal muscle enzymes and blue urine in a farmworker from Thailand. *Isr Med Assoc J* 2001;**3**:704.
98. SHIOTA T, ARIZONO N, YOSHIOKA T et al. Imported trichinellosis with severe myositis—report of a case. *Kansenshogaku Zasshi* 1999;**73**:76–82.
99. GROSS B, OCHOA J. Trichinosis: clinical report and histochemistry of muscle. *Muscle Nerve* 1979;**2**:394–8.
100. SUNDARAY NK, DAS AK, SANCHETEE PC, ANAND AC, SINGH W. Acute filarial myositis. *J Assoc Physicians India* 1992;**40**:693–4.
101. HUANG BK, MONU JU, DOUMANIAN J. Diabetic myopathy: MRI patterns and current trends. *AJR Am J Roentgenol* 2010;**195**:198–204.
102. SU Z, ROBINSON A, HU L et al. Acupuncture plus Low-Frequency Electrical Stimulation (Acu-LFES) Attenuates Diabetic Myopathy by Enhancing Muscle Regeneration. *PLoS ONE* 2015;**10**:e0134511.
103. D'SOUZA DM, AL-SAJEE D, HAWKE TJ. Diabetic myopathy: impact of diabetes mellitus on skeletal muscle progenitor cells. *Front Physiol* 2013;**4**:379. doi:10.3389/fphys.2013.00379.
104. MAZOCH MJ, BAJAJ G, NICHOLAS R et al. Diabetic myonecrosis: likely an underrecognized entity. *Orthopedics* 2014;**37**:e936–9.
105. CHALASANI S, BETTADAHALLI SS, BHUPATHI SV, ASWANI VH. A novel case of diabetic muscle necrosis in a patient with cystic fibrosis-related diabetes. *Clin Med Res* 2013;**11**:113–6.
106. YU CP, LIU TH, WANG B, ZHANG Q. Diabetic muscle infarction in a patient admitted to the cardiac ward. *Intern Med* 2014;**53**:1227–9.
107. MADHU SV, JAIN R, KANT S, PRAKASH V, KUMAR V. Myopathy presenting as a sole manifestation of hypothyroidism. *J Assoc Physicians India* 2010;**58**:569–70.
108. MATAYOSHI T, OMI T, MAYUMI N, KAWANA S. Hashimoto's thyroiditis with clinical manifestations resembling dermatomyositis: a case report. *J Nippon Med Sch* 2014;**81**:106–9.
109. SHAHEEN D, KIM CS. Myositis associated with the decline of thyroid hormone levels in thyrotoxicosis: a syndrome? *Thyroid* 2009;**19**:1413–7.
110. KIM JM, SONG EJ, SEO JS, NAM EJ, KANG YM. Polymyositis-like syndrome caused by hypothyroidism, presenting as camptocormia. *Rheumatol Int* 2009;**29**:339–42.
111. FARIAS MOELLER R, ZECAVATI N, SHERAFAT-KAZEMZADEH R, ALEINIKOFF S, RENNERT W. Adolescent with rhabdomyolysis due to undiagnosed hypothyroidism. *Case Rep Pediatr* 2011;**201**:670673.
112. TACHIBANA S, YOKOI T, SATO S, ODA Y, YANASE T, YAMASHITA H. Idiopathic orbital myositis associated with Graves' disease. *Intern Med* 2013;**52**:787–90.
113. HARDIMAN O, MOLLOY F, BRETT F, FARRELL M. Inflammatory myopathy in thyrotoxicosis. *Neurology* 1997;**48**:339–41.
114. WALKER RP, PALOYAN E, GOPALSAMI C. Symptoms in patients with primary hyperparathyroidism: muscle weakness or sleepiness. *Endocr Pract* 2004;**10**:404–8.
115. THAGE O. Metabolic neuropathies and myopathies in adults. Clinical aspects. *Acta Neurol Scand* 1970;**46** (Suppl 43):120.
116. AKMAL M. Rhabdomyolysis in a patient with hypocalcemia due to hypoparathyroidism. *Am J Nephrol* 1993;**13**:61–3.
117. MUNOZ J, FERRARI N, KURIAKOSE P. Iron-overload myopathy. *Int J Hematol* 2011;**94**:503–4.
118. BREGMAN H, GELFAND MC, WINCHESTER JF, MANZ HJ, KNEPESHIELD JH, SCHREINER GE. Iron-overload-associated myopathy in patients on maintenance haemodialysis: a histocompatibility-linked disorder. *Lancet* 1980;**2**:882–5.
119. WOUTHUIS SF, VAN DEURSEN CT, TE LINTELO MP, ROZEMAN CA, BEEKMAN R. Neuromuscular manifestations in

- hereditary haemochromatosis. *J Neurol* 2010;**257**:1465–72.
120. MOHAMMAD A, CAREY JJ, STORAN E, SCARRY M, COUGHLAN RJ, LEE JM. High prevalence of fibromyalgia in patients with HFE-related hereditary hemochromatosis. *J Clin Gastroenterol* 2013;**47**:559–64.
 121. SWINKELS DW, AALBERS N, ELVING LD, BLEIJENBERG G, SWANINK CM, VAN DER MEER JW. Primary haemochromatosis: a missed cause of chronic fatigue syndrome? *Neth J Med* 2002;**60**:429–33.
 122. MANOLI I, KWAN JY, WANG Q et al. Chronic myopathy due to immunoglobulin light chain amyloidosis. *Mol Genet Metab* 2013;**108**:249–54.
 123. JEON CY, JIN SY, JANG JW, KIM US. A case of isolated primary amyloidosis in the medial rectus muscle. *Graefes Arch Clin Exp Ophthalmol* 2013;**251**:395–6.
 124. CARR AS, PELAYO-NEGRO AL, JAUNMUKTANE Z et al. Transthyretin V122I amyloidosis with clinical and histological evidence of amyloid neuropathy and myopathy. *Neuromuscul Disord* 2015;**25**:511–5.
 125. OHTSUKA Y, YASUI N, SEKIGUCHI K et al. Case of amyloidosis with amyloid deposition detected only in skeletal muscles. *Rinsho Shinkeigaku (Clin Neurol)* 2012;**52**:739–43.
 126. COLOMBO I, FRUGUGLIETTI ME, NAPOLI L et al. IgD Multiple Myeloma Paraproteinemia as a Cause of Myositis. *Neurol Res Int* 2010;**2010**:808474.
 127. MATSUMURA E, YAMAGUCHI T, TOMIDOKORO Y, ISHII A, TAMAOKA A. A representative case of joint contracture as a main feature of AL amyloid deposits identified in the skeletal muscles. *Rinsho Shinkeigaku (Clin Neurol)* 2014;**54**:907–10.
 128. OSTROW LW, CORSE AM, MORRISON BM et al. Expanding the spectrum of monoclonal light chain deposition disease in muscle. *Muscle Nerve* 2012;**45**:755–61.
 129. ABUZINADAH AR, JOSEPH JT, KORNGUT L. Amyloid myoneuropathy mimicking inclusion body myositis. *Can J Neurol Sci* 2013;**40**:255–8.
 130. IKEZAWA Y, OKA K, NAGAYAMA R et al. Bence-Jones protein-type myeloma with amyloid myopathy presenting as amyloidomas and extensive amyloid deposits in the muscularis propria: a rapidly fatal autopsy case. *Int J Surg Pathol* 2012;**20**:83–8.
 131. LAHORIA R, WINDER TL, LUI J, AL-OWAIN MA, MILONE M. Novel ANO5 homozygous microdeletion causing myalgia and unprovoked rhabdomyolysis in an Arabic man. *Muscle Nerve* 2014;**50**:610–3.
 132. MARSDEN JT, PETERS TJ. Rhabdomyolysis in a patient with acute intermittent porphyria. *Ann Clin Biochem* 2004;**41**:341–3.
 133. POERSCH M. Acute intermittent porphyria with transient paresis and contracture. Evidence for initial myopathic dysfunction? *Nervenarzt* 1998;**69**:171–3.
 134. COHEN PL, HADLER NM, STARKENBURG R. Acute intermittent porphyria presenting as acute muscle pain, fever, and weakness. *Arthritis Rheum* 1997;**40**:586–7.
 135. KAYA A, KARA M, TIFTIK T et al. Ultrasonographic evaluation of the muscle architecture in patients with systemic lupus erythematosus. *Clin Rheumatol* 2013;**32**:1155–60.
 136. SCHUBERT J, MANSCHWETUS H, WALTER GF, ZEIDLER H, SCHMIDT RE. Systemic lupus erythematosus (SLE) with panniculitis and rhabdomyolysis. *Immun Infekt* 1991;**19**:58–9.
 137. JAKATI S, RAJASEKHAR L, UPPIN M, CHALLA S. SLE myopathy: a clinicopathological study. *Int J Rheum Dis* 2015;**18**:886–91.
 138. ENDO Y, IKAWA M, TAKAHASHI N, NISHINO I, SUZUKI S, YONEDA M. A case of systemic lupus erythematosus predominantly presenting with myositis at onset. *Rinsho Shinkeigaku (Clin Neurol)* 2013;**53**:634–7.
 139. SANTOSA A, VASOO S. Orbital myositis as manifestation of systemic lupus erythematosus—a case report. *Postgrad Med J* 2013;**89**:59.
 140. GARCÍA-REYNOSO MJ, VERAMENDI-ESPINOZA LE, RUIZ-GARCIA HJ. Ascending paresis as presentation of an unusual association between necrotizing autoimmune myopathy and systemic lupus erythematosus. *Reumatol Clin* 2014;**10**:183–6.
 141. SL, TONY K, RAGHUPATHY VS, MALEPATI B. A rare case of mixed connective tissue disease (MCTD) with intricate features of lupus, polymyositis and rheumatoid arthritis presenting with severe myositis. *J Clin Diagn Res* 2015;**9**:OD05–7.
 142. KIJIMA T, KANEKIYO S, NAKASUGA C et al. A case report of a patient with overlap syndrome systemic lupus erythematosus(SLE) and polymyositis (PM) whose condition improved following treatment for coexisting descending colon cancer. *Gan To Kagaku Ryoho* 2013;**40**:1936–8.
 143. DE CARVALHO JF, DA MOTA LM, BONFA E. Fatal rhabdomyolysis in systemic lupus erythematosus. *Rheumatol Int* 2011;**31**:1243–5.
 144. COLAFRANCESCO S, PRIORI R, GATTAMELATA A et al. Myositis in primary Sjögren's syndrome: data from a multicentre cohort. *Clin Exp Rheumatol* 2015;**33**:457–64.
 145. LIN DF, YAN SM, ZHAO Y et al. Clinical and prognostic characteristics of 573 cases of primary Sjögren's syndrome. *Chin Med J (Engl)* 2010;**123**:3252–7.
 146. DANLOS FX, DAOUED F, SEROR R, MARIETTE X. Orbital Myositis and Primary Sjögren Syndrome. *J Rheumatol* 2015;**42**:1536–7.
 147. HUANG CT, CHEN YC, LIN C, HSIAO YC, SHEU LF, TU MC. A case of dermatomyositis with secondary Sjögren's Syndrome-diagnosis with follow-up study of technetium-99 m pyrophosphate scintigraphy. *Acta Neurol Taiwan* 2013;**22**:162–8.
 148. CARRON M, IEPPARIELLO G, ORI C. Sugammadex in a patient with Sjogren's syndrome and polymyositis. *Br J Anaesth* 2013;**111**:1034–5.
 149. MISTERSKA-SKÓRA M, SEBASTIAN A, DZIĘGIEL P, SEBASTIAN M, WILAND P. Inclusion body myositis associated with Sjögren's syndrome. *Rheumatol Int* 2013;**33**:3083–6.
 150. ROJANA-UDOMSART A, NEEDHAM M, LUO YB et al. The association of sporadic inclusion body myositis and Sjögren's syndrome in carriers of HLA-DR3 and the 8.1 MHC ancestral haplotype. *Clin Neurol Neurosurg* 2011;**113**:559–63.
 151. ANCUȚA C, POMIRLEANU DC, ANTON CR et al. Rheumatoid myositis, myth or reality? A clinical, imaging and histological study. *Rom J Morphol Embryol* 2014;**55**:781–5.
 152. JO SE, KIM YJ, LEE KH et al. Muscular involvement of Behçet's disease: ultrasonography, computed tomography, and magnetic resonance imaging findings. *Clin Imaging* 2012;**36**:643–6.
 153. COUDERC M, MATHIEU S, TOURNADRE A, DUBOST JJ, SOUBRIER M. Acute ocular myositis occurring under etanercept for rheumatoid arthritis. *Joint Bone Spine* 2014;**81**:445–6.
 154. FADINI GP. A case of orbital myositis associated with rheumatoid arthritis. *Ann Rheum Dis* 2003;**62**:383.
 155. MEYER A, LANNES B, CARAPITO R et al. Eosinophilic myositis as first manifestation in a patient with type 2

- myotonic dystrophy CCTG expansion mutation and rheumatoid arthritis. *Neuromuscul Disord* 2015;**25**:149–52.
156. LÓRÁND V, CZIRJÁK L, MINIER T. Musculoskeletal involvement in systemic sclerosis. *Presse Med* 2014;**43**:e315–28.
 157. VEMULAPALLI S, SHARER LR, HSU VM. Inclusion body myositis in a patient with RNA polymerase III antibody-positive systemic sclerosis. *J Rheumatol* 2015;**42**:730–2.
 158. JUNG M, BONNER A, HUDSON M, BARON M. Pope JE; Canadian Scleroderma Research Group (CSRG). Myopathy is a poor prognostic feature in systemic sclerosis: results from the Canadian Scleroderma Research Group (CSRG) cohort. *Scand J Rheumatol* 2014;**43**:217–20.
 159. FERNÁNDEZ-SERNA M, ARBOLEYA L, ALONSO S, QUEIRO R, ALPERI M. Dropped head syndrome in a patient with scleromyositis. *J Clin Rheumatol* 2013;**19**:32–4.
 160. SCHANZ S, HENES J, ULMER A et al. Magnetic resonance imaging findings in patients with systemic scleroderma and musculoskeletal symptoms. *Eur Radiol* 2013;**23**:212–21.
 161. MAUHIN W, RIVIÈRE S, CABANE J, TIEV KP. Improvement in lung fibrosis using intravenous immunoglobulin in systemic sclerosis with myositis. *Scand J Rheumatol* 2014;**43**:170–1.
 162. LIEWLUCK T, ERNSTE FC, TRACY JA. Frequency and spectrum of myopathies in patients with psoriasis. *Muscle Nerve* 2013;**48**:716–21.
 163. SSI-YAN-KAI I, PEARSON A. Orbital myositis and psoriatic arthritis. *Can J Ophthalmol* 2012;**47**:e42–3.
 164. BAINS S, REIMERT M, WIN AZ, KHAN S, APARICI CM. A Patient with Psoriatic Arthritis Imaged with FDG-PET/CT Demonstrated an Unusual Imaging Pattern with Muscle and Fascia Involvement: A Case Report. *Nucl Med Mol Imaging* 2012;**46**:138–43.
 165. MACHADO NP, CAMARGO CZ, OLIVEIRA AC, BUOSI AL, PUCINELLI ML, SOUZA AW. Association of anti-glomerular basement membrane antibody disease with dermatomyositis and psoriasis: case report. *Sao Paulo Med J* 2010;**128**:306–8.
 166. ALLALI D, SEEBACH JD. The anti-synthetases syndrome: diagnostic and Treatments. *Rev Med Suisse* 2015;**11**:808–12.
 167. DIEVAL C, DELIGNY C, MEYER A et al. Myocarditis in Patients With Antisynthetase Syndrome: Prevalence, Presentation, and Outcomes. *Medicine (Baltimore)* 2015;**94**:e798.
 168. JOUNEAU S, HERVIER B, JUTANT EM et al. Pulmonary manifestations of antisynthetase syndrome *Rev MalRespir* 2015;**32**:618–28.
 169. EL-HATTAB AW, ADESINA AM, JONES J, SCAGLIA F. MELAS syndrome: Clinical manifestations, pathogenesis, and treatment options. *Mol Genet Metab* 2015;**116**:4–12.
 170. CONWAY R, BERGIN D, CAREY JJ, COUGHLAN RJ. A case of Behcet's syndrome presenting as focal myositis. *Rheumatology (Oxford)* 2012;**51**:975.
 171. CHEBBI W, JERBI S, AMMARI W, YOUNES M, SFAR MH. Orbital myositis in Behçet's disease. *Joint Bone Spine* 2014;**81**:264.
 172. DURSUN D, AKOVA Y, YÜCEL E. Myositis and scleritis associated with Behçet's disease: an atypical presentation. *Ocul Immunol Inflamm* 2004;**12**:329–32.
 173. OGOSE T, TAMAKI W, SHINAHARA K, KANEKO M, WAKATA Y, NARUSE K. A case of recurrent myositis as the main manifestation of Behçet disease. *Pediatr Int* 2010;**52**:e101–4.
 174. ABADI U, HADARY R, SHILO L, SHABUN A, GREENBERG G, KOVATZ S. Myositis in Behçet's disease after tonsillectomy. *Isr Med Assoc J* 2006;**8**:294–5.
 175. SARUI H, MARUYAMA T, ITO I et al. Necrotising myositis in Behçet's disease: characteristic features on magnetic resonance imaging and a review of the literature. *Ann Rheum Dis* 2002;**61**:751–2.
 176. LINGENFELSER T, DUERK H, STEVENS A, GROSSMANN T, KNORR M, SAAL JG. Generalized myositis in Behçet disease: treatment with cyclosporine. *Ann Intern Med* 1992;**116**:651–3.
 177. SALAM A, MELIGONIS G, MALHOTRA R. Superior oblique myositis as an early feature of orbital Wegener's granulomatosis. *Orbit* 2008;**27**:203–6.
 178. MANTEGANI A, AUBERT J, GUELPA G. Inflammatory myopathy: unusual presentation of a case of Wegener disease. *Schweiz Med Wochenschr* 1993;**123**:1996–2000.
 179. PARENT ME, LARUE S, ELLEZAM B. Eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome) presenting as diffuse myositis. *BMC Musculoskelet Disord* 2014;**15**:388.
 180. LAZZARIN P, PRESOTTO F, POLO A. Churg-Strauss syndrome presenting as polymyositis: report of a case. *Reumatismo* 2009;**61**:140–4.
 181. FUJII T, NORIZUKI M, KOBAYASHI T, YAMAMOTO M, KISHIMOTO M. A case of eosinophilic orbital myositis associated with CSS. *Mod Rheumatol* 2010;**20**:196–9.
 182. CONNES P. Sickle cell trait, exertional rhabdomyolysis, and compartment syndrome. *Lancet* 2015;**385**:1948.
 183. SHRIVASTAV A, KUMAR V, PAL J. Dermatomyositis associated with acute myelocytic leukemia. *Rheumatol Int* 2010;**30**:671–3.
 184. ALJOHANI NI, CARETTE S, LIPTON JH. Inclusion body myositis in a patient with chronic myeloid leukemia treated with dasatinib: a case report. *J Med Case Rep* 2015;**9**:214.
 185. BECK EH, AMATO AA, GREENBERG SA. Inclusion body myositis and chronic lymphocytic leukemia: a case series. *Neurology* 2014;**83**:98–9.
 186. CHEN MC, YANG SH, YAO TK, CHONG PN, CHEN SH. Bilateral hip pain caused by adductor pyomyositis as the initial presentation of chronic myeloid leukemia in a 17-year-old child. *Pediatr Neonatol* 2011;**52**:353–7.
 187. TRAINA ED, CRIDER MF, BERMAN D, SPOTO-CANNONS A. Pyomyositis as the presenting feature of acute lymphocytic leukemia. *Clin Pediatr (Phila)* 2008;**47**:167–70.
 188. CHEN YC, SHEEN JM, HUANG LT, WU KS, HSIAO CC. Disseminated tuberculous myositis in a child with acute myelogenous leukemia. *Pediatr Neonatol* 2009;**50**:74–7.
 189. FUKUSHIMA T, IWAO H, NAKAZIMA A et al. MRSA-pyomyositis in a patient with acute myelogenous leukemia after intensive chemotherapy. *Anticancer Res* 2009;**29**:3361–4.
 190. PARK JK, SUBHAWONG AP, ZIMINSKI CM, GRADER-BECK T. Dermatomyositis as the initial presentation of a large anaplastic T-cell lymphoma. *J Clin Oncol* 2011;**29**:e378–80.
 191. MARIE I, GUILLEVIN L, MENARD JF et al. Hematological malignancy associated with polymyositis and dermatomyositis. *Autoimmun Rev* 2012;**11**:615–20.
 192. DIAS JS, DITTRICH MA, SOUSA JM, TEIXEIRA LF, MANSO PG. Medial rectus muscle myositis as an atypical presentation of mucosa-associated lymphatic tissue lymphoma: a case report. *Arq Bras Oftalmol* 2014;**77**:116–8.

Finsterer et al.

193. FRIMAN G. Serum creatine phosphokinase in epidemic influenza. *Scand J Infect Dis* 1976;**8**:13–20.
194. SHEPHERD JJ. Tropical myositis: is it an entity and what is its cause? *Lancet* 1983;**2**:1240–2.
195. LINDBERG C, ANDERSON B, ENGVALL M, HULT M, OLDFORS A. Search for Pompe disease among patients with undetermined myopathies. *Acta Neurol Scand* 2015. DOI: 10.1111/ane.12460. [Epub ahead of print].
196. SWASH M, SCHWARTZ MS, VANDENBURG MJ, POLLOCK DJ. Myopathy in Whipple's disease. *Gut* 1977;**18**:800–4.
197. HENRIKSSON KG, HALLERT C, NORRBY K, WALAN A. Polymyositis and adult coeliac disease. *Acta Neurol Scand* 1982;**65**:301–19.
198. LI X, MAO YT, WU JJ, LI LX, CHEN XJ. Anti-AMPA receptor encephalitis associated with thymomatous myasthenia gravis. *J Neuroimmunol* 2015;**281**: 35–7.
199. GLUECK CJ, CONRAD B. Severe vitamin d deficiency, myopathy, and rhabdomyolysis. *N Am J Med Sci* 2013;**5**:494–5.